Lecture 25: Sub-Surface and Volume Scattering

Yuhao Zhu

http://yuhaozhu.com yzhu@rochester.edu CSC 259/459, Fall 2025 Computer Imaging & Graphics

The Roadmap

Theoretical Preliminaries


Human Visual Systems

Display and Camera

Modeling and Rendering

3D Modeling
Rasterization and GPU
Ray Tracing
Shading
Rendering in AR/VR

Subsurface/Volume Scattering



So far we consider only surface reflection, but not how photons interact with particles under surface.


Particles of translucent materials under the material surface scatter lights like crazy and some scattered photons make their way out of the surface.


Material Types


Pure absorption (Transparent)


A little scattering + absorption (Translucent)

A lot of scattering + absorption (Opaque)














Volume Scattering

Particles scatter and absorb photons, and it's important to model in:

- Translucent materials (e.g., jade, skin), which have strong subsurface scattering (SSS) in addition to surface scattering. SSS can be ignored in cases where refraction is weak or particle absorption is strong so that the back-scattering doesn't contribute much to the overall material appearance.
- "Participating media" (e.g., cloud, smoke), which have no surface and everything in volume scattering.







https://assetstore.unity.com/packages/vfx/shaders/jade-translucency-subsurface-scattering-shader-84486




General Idea


The principle way to model volume scattering is the Radiative Transfer theory, which is an integro-differential equation that generally have no analytical solution.

We solve it by:


- make simplified assumptions (e.g., Kubelka-Munk model for diffuse illumination and isotropic scattering, diffusion approximation applied to BSSDF [Jensen et al. 2000])
- plus: using numerical methods to approximate the solution (e.g, monte-carlo estimator)

Volume Scattering Processes






Volume Scattering Processes




Pure Absorption

Modeling Pure Absorption


Consider a very thin slice (so thin that no particles are covering each other)





Modeling Pure Absorption

Consider a very thin slice (so thin that no particles are covering each other)




Percentage of absorbed photons

$$\frac{\rho E \Delta s A}{E} = \frac{\Delta I}{I}$$

$$\frac{\Delta I}{\Delta s} = -\rho IA$$

Beer-Lambert Law



As s increases (light travel farther), the intensity decreases exponentially!

$$\frac{dI}{ds} = -\rho IA$$


$$\int V$$


$$I = I_0 e^{-\rho As}$$
 If density is uniform
$$I = I_0 e^{-\int_0^s \rho(s)A}$$
 If density is non-uniform

"The person credited with a scientific law is almost certainly not the one who discovered it."

<u>Bouguer</u> was the first to discover this law.


Beer-Lambert Law





Exponential Decay

$$I = I_0 e^{-\rho As}$$



Exponential decay happens when a quantity decreases at a rate proportional to its current value.


• Opposite of exponential growth.

Exponential decay means lights are hard to be completely absorbed.

• 10% of light @ 70 m deep under ocean, but you can still see lights at 850 m deep.

Many real-world scenarios exhibit exponential growth/decay.


Exponential Growth



Source: Post reporting from state health department data

Every exponential growth will come to an end when...






New York Times

Absorption Efficiency and Coefficient

Density/concentration Cross-section (Unit: 1/m³) area (Unit: m²)

$$I = I_0 e^{-\rho As}$$

More generally



Absorption efficiency (unitless): the probability that a photon is actually absorbed when it hits a particle.

$$I = I_0 e^{-\rho Q_a As}$$

Absorption coefficient:

Unit: 1/m

$$\mu = \rho Q_a A$$

Percentage of light left

$$I = I_0 e^{-\mu s} \qquad \frac{I}{I_0} = e^{-\mu s}$$

* As always, absorption efficiency and coefficient are wavelength-dependent.


$$C_a = Q_a A$$
 is sometimes called effective absorption cross section

Absorption Efficiency and Coefficient

Informally, μ quantifies how much light is absorbed per unit distance.

- $\mu = 1$ means after passing through 1 m the radiance is reduced by a factor of e.
- Another way to understand μ is that $\mu = 0.1$ means a photon can travel about 10 m before being absorbed. (1/ μ is the mean lifetime in an exponential decay).

Photoreceptor Absorption




Percentage of light left
$$\frac{I}{I_0} = e^{-\mu s}$$

Outer segment is light-sensitive part of rods. It's about 25 μ m long.

- It absorbs about 3% of the incident light for every µm.
- How much light is absorbed by a rod cell? (Answer 53%)

Absorption/Transmission vs. Concentration

$$I = I_0 e^{-\rho Q_a A s}$$



Transmittance (1 - absorption) of three colored-ink at various concentrations (\mathbf{c}).

Pure Scattering

(from a single particle)

Modeling Scattering

We need to model two aspects of scattering:

- How much is light scattered. This is captured by scattering coefficient.
- In which direction is a photon scattered. This is is captured by the so-called "phase function".

The general way to model scattering behavior amounts to solving Maxwell's equations for scattering. It's called the Mie solution.

• "Gustav Mie is the last person to discover the scattering behavior that bears his name."

Importantly, people have worked out approximations under different conditions, using which we can readily explain many scattering behaviors.

Modeling Scattering

We will first consider cases where a photon is scattered by one particle.


- This happens when the particles are reasonably apart of each other.
- A photon can be scattered by multiple particles if the particle concentration is high or when the material is thick.

We will then see general cases of multiple scattering + absorption.

We also assume elastic scattering, where scattered photon has the same wavelength as the incident photon.


• There are inelastic scattering, which does happen in nature but we won't consider here.

Modeling Scattering



^{*} even though here we assume a photon is scattered by only a single particle, multiple particles can be scattering different photons.

Scattering efficiency (unitless): the probability that a photon is actually back-scattered when it hits a particle.



$$\mu = \rho Q_{S}A$$
 is called scattering coefficient: Unit: 1/m

$$C_S = Q_S A$$
 is called effective scattering cross section


Three Scattering Regimes


Small particles (relative size $< 0.1 \lambda$) Rayleigh regime

Medium particles $[0.1 \lambda, 100 \lambda]$ Mie regime

Large particles (relative size $> 100 \lambda$) **Geometry** regime

Soot particle scattering 500 nm photons






Three Scattering Regimes

Small particles (relative size $< 0.1 \lambda$) Rayleigh regime

Medium particles $[0.1 \lambda, 100 \lambda]$ Mie regime

Large particles (relative size $> 100 \lambda$) **Geometry** regime

Water drop scattering 500 nm photons





General Trends


Scattering efficiency

• Small particles scatter little; large particles scatter a lot.


Scattering direction

- Small particles scatter lights both forward and backward.
- More forward direction as particle size increases.

Q_s: Scattering efficiency



Qf: Forward scattering efficiency. Q_f/Q_s shows how much light is scattered forward.



Spherical gold particle scattering 500 nm photons.

As particle size increases

• significantly more forward scattering + higher scattering efficiency


(d) (e) (b) (a) 0.1 1.0 particle radius = 50 nm 10 nm 100 nm particle radius = 500 nm 1000 nm $\sim 0.18 \lambda$ $\sim 0.91 \lambda$ $\sim 1.82 \lambda$ $\sim 0.02 \lambda$ $\sim 0.1 \lambda$

Scattering 550 nm photons.

As particle size increases

• significantly more forward scattering + higher scattering efficiency

Length is proportional to scattering efficiency



Small Large

Small cross section area

Low scattering efficiency

Even backward and forward scattering

Large cross section area


High scattering efficiency

Mostly forward scattering

Increase transparency Length is proportional to scattering efficiency

Even backward and

forward scattering




Small Large Large cross section area Small cross section area Low scattering efficiency High scattering efficiency Increase opacity

Colour and the Optical Properties of Materials 3ed, Tilley 34

Mostly forward scattering

Length is proportional to scattering efficiency

"Hiding Power" vs. Size



$$\frac{H}{V} = Q_s A(1-g) \propto \frac{Q_s(1-g)}{r}$$

There is a sweet spot in particle size that provides the highest opacity (hiding light behind it).

• Accidentally the sweet spot falls roughly within the visible range.


Hippopodius hippopus is normally transparent, but when disturbed its protein precipitate and form larger particles. It then becomes opaque.



Impact of Refractive Index

Part of scattering is caused by the refractive index mismatch between the particle and the surrounding medium.


- Water droplets surrounded by air in fog; pigments surrounded by oil in paint, etc.
- It's the relative index, **m**, between the particle and the medium that matters.



$$m = rac{n_{particle}}{n_{medium}}$$
 $N_{air} \approx 1$
 $N_{water} \approx 1.33$
 $N_{glass} \approx 1.51$
 $N_{SiO2} \approx 1.46$

Aside: How Paints are Made

Paint ~= Pigment + Binder




Binder (medium/vehicle) provides the adhesion needed to hold the pigments together.

Usually some additives are added:

- Glaze
- Extender to increase durability
- Glitter for shimmer

Paint ~= Pigment + Binder




The incident lights are also reflected straight away at the air-medium surface (dashed lines)

 We will ignore this for now and come back to it later

Note that surface reflection is just the macroscopic behavior of photon scattering off of the binder particles (scattering with interference).


Paint on Substrate





https://feltmagnet.com/painting/How-To-Choose-A-Canvas-For-Painting 41


Paint on Substrate



The paint is applied to a substrate, e.g., a canvas or paper. The substrate is nothing more than another layer of scattering/ absorbing material.


 We will show how layered material is modeled later.

Mona Lisa, Leonardo Da Vinci, 1506. The Louvre.







The Birth of Venus, Sandro Botticelli, c 1485. Uffizi Galley, Florence.



Portrait of a Young Woman in Red, A.D. 90–120. The Met.




Mummy Portrait of a Woman, A.D. 100. Getty.




Keep Moving Forward, Darien Bogart






Stephen Quiller



The Blue Boat, Winslow Homer, 1892. Museum





"Watercolor" painting. Binder is Gum Arabic + water.

Pigment Size

average pigment particle size				
microns	meters	representative pigments		
1000µm	10 ⁻³	= 1 millimeter		
500µm				
100µm	10 ⁻⁴	coarse historical mineral pigments modern luster & iridescent pigments		
50μm		smallest particles visible without magnification cobalt violet manganese blue		
10µm	10 ⁻⁵	cobalt green cobalt turquoise cerulean blue manganese violet black iron oxides		
5µm		ultramarine blue (RS) viridian cobalt blue violet (brown) iron oxides yellow iron oxides		

1µm	10 ⁻⁶	ultramarine blue (GS) red iron oxides cadmium red cadmium orange semiopaque synthetic organics diarylides pyrroles naphthols perinone orange
0.5μm		= 500 nanometers = wavelength of "blue green" light chromium oxide green cadmium yellow bismuth yellow titanium white transparent red iron oxides transparent yellow iron oxides semitransparent synthetic organics arylides benzimidazolones dioxazines perylenes
0.1µm	10 ⁻⁷	zinc white iron [prussian] blue transparent synthetic organics quinacridones phthalocyanines
0.05µm		carbon black
0.01µm	10 ⁻⁸	
0.0015µm	10 ⁻⁹	glucose (corn syrup) molecule
0.0003µm	10 ⁻¹⁰	water molecule

Disperse Pigments



Pigments are very tiny but usually agglomerate, which is the powders we buy.

We need to separate the pigment particles from the agglomerates and disperse them into the binder.








Back From Aside

Refractive Index vs. Scattering Efficiency

$$m = \frac{n_{particle}}{n_{medium}}$$

wavelength = 500 nm




Observations:

- Large relative refractive index (m) increases scattering efficiency
- When m is close to 1, very little scattering until closer to the geometric (large) regime
- The scattering efficiency in the geometric regime is independent of refractive index *m*,

Refractive Index vs. Scattering Direction

As m approaches 1 (i.e., the refractive indices of the medium and particle become more similar), forward scattering is more significant.


• Extreme case is when the medium and the particle are the same material, then scattering is completely forward ($\theta_1 = \theta_2$)!




Why Are Wet Things Darker?

$$N_{SiO2} \approx 1.46$$









Why Are Wet Things Darker?


The RI of sand is much closer to water than to air. So light scattered forward more and travels longer distance and has more opportunities to be absorbed.

Note that scattering by sand is in the geometric regime (typical sand size is [62.5 μ m, 2 mm]), so scattering efficiency is high regardless of RI. The question is where photons scatter.






The Miracles of Saint Francis of Paola, P. P. Rubens, 1627–8. J. Paul Getty Museum.



The refractive index of the linseed oil in the paint increases over time and matches that of the pigment. The paint layer becomes transparent, revealing the pentimento underneath.



Aside: Pentimento

The presence/emergence of earlier paints that have been painted over.

- Painter changing mind (pentimento is from "pentirsi" in Italian, which means repentance)
- Underdrawing

Can be identified using imaging techniques (e.g., X-ray), or from paint composition changes (previous slide).

Portrait of Pope Julius II, Raphael, 1511-1512. National Gallery, London.



The Old Guitarist, Pablo Picasso, 1903-1904. The Art Institute of Chicago.





Acrylic Paints Dry Dark

Wet Dry

Naphthol Red








Acrylic Paints Dry Dark

Wet Dry




Naphthol Red

 $N_{naphthol\ reds} \approx 1.60$



Ultramarine Blue

 $N_{ultramarine-blue} \approx 1.50$



 $N_{water} \approx 1.33$ $N_{acrylic\ binder} \approx 1.49$

Reason: when water evaporates, the refractive indices of pigments and binder are closer, which encourages more forward scattering, which increases total absorption.

General Scenarios

(Absorption + Multiple Scattering)

Multiple Scattering

So far we assume that a photon is either absorbed or scattered by at most one particle before entering our eyes.

 Photon scattered by a particle will find its way out without encountering another scattering particle.

With high particle concentration or a thick colorant layer, photons can be scattered by multiple particles and might later be absorbed.

• A photon scattered might be later re-scattered.

Multiple scattering + absorption is the common case in nature, arts, and tech.


Solutions

Our goal is to estimate the (spectral) **reflectance** of a material (colorant) in the presence of multiple scattering and absorption.

There are two approaches in general:

- The general case amounts of solving the so-called **Radiative Transfer Equations**, for which analytical solutions are hard to obtained.
- Again, we make simplifications to allow modeling special, but common cases.

Radiative Transfer Equations




Intuition: we account for the change of radiance at p along direction ω by

- Removing the absorbed and out-scattered (extinction) portion of incident ray
- Adding emission
- Adding in-scattering (integrated over all directions)


Phase function: radiance distribution

BSSDF: Combining Surface and Volume Scattering

Bidirectional Scattering Surface Reflectance Distribution Function (BSSRDF).



Integrate over all points on the surface and from all directions



Models how the irradiance of incident rays affects






Perfect Diffuse Model

The assumptions we make are:

- Particles have the same scattering and absorption coefficients (homogeneity)
- Particles are randomly oriented (special case: they are all spheres)
- Incident light is uniform at all directions at all surface points (diffuse illumination)

Perfect Diffuse Model



Total downward flux

Total upward flux

Given the assumptions, there is no difference between different horizontal points (**Y**) at the same depth (**X**): the downward (and upward) flux for all **Y**s at the same **X** is the same.


So the reflectance of any point is equivalent to the ratio of the total downward flux and the total upward flux at the depth where the point is.

Kubelka-Munk Equation

K: absorption coefficient

S: back-scattering coefficient

X: thickness



Total downward flux at the top of the layer


Total upward flux at the bottom of the layer


Kubelka-Munk Equation

K: absorption coefficient

S: back-scattering coefficient

X: thickness





Boundary conditions:

$$R_g = \frac{j_0}{i_0}$$


Reflectance of the layer:

$$R_X = \frac{j_X}{i_X}$$

This is what we want to calculate!

Kubelka-Munk Equation: General Solution

$$R_X = \frac{1 - R_g[a - b \coth(bSX)]}{a - R_g + b \coth(bSX)} \qquad a = \frac{K + S}{S} \qquad b = \sqrt{a^2 - 1} \qquad \coth x = \frac{e^{2x} + 1}{e^{2x} - 1}$$




Reflectance depends on:

- Layer thickness X
- Absorption and scattering coefficients K and S
- Background reflectance: R_g


Kubelka-Munk Equation: No Scattering

$$S=0$$



% of photons left after they reach the bottom


% of photons left after they reach the top again



% of photons reflected back by the substrate

Kubelka-Munk Equation: No Substrate

$$R_g = 0$$
 $R_X = \frac{1}{a + b \coth(bSX)}$ $a = \frac{K + S}{S}$ $b = \sqrt{a^2 - 1}$




Substrate reflectance $\mathbf{R_g}$ is 0 now. Alternatively this can thought of as having a black substrate. The effect is the same: whatever gets out of the layer never comes back.

Kubelka-Munk Equation: Complete Hiding

$$\lim_{X \to \infty} R_X = R_{\infty} = \frac{1}{a+b} = 1 + \frac{K}{S} - \sqrt{\left(\frac{K}{S}\right)^2 + 2\frac{K}{S}}$$


$$\frac{K}{S} = \frac{(1 - R_{\infty})^2}{2R_{\infty}}$$



When the layer is so thick that the background reflectance is irrelevant, the resulting reflectance depends only on the absorption and scattering coefficients.

We say the "hiding is complete".

Hiding Power in Painting



$$HP = \frac{R_{black}}{R_{white}} \begin{tabular}{ll} Reflectance when substrate is black. \\ Reflectance when substrate is white. \\ Reflec$$

In paint industry, the way they quantify hiding power is to see how the surface reflectance changes when the background is white vs. black.


If the layer's hiding is complete, the reflectance ratio should 1, i.e., the background color doesn't matter and the colorant is **opaque**.


Reflectance vs. Concentration

$$\frac{K}{S} = \frac{(1 - R_{\infty})^2}{2R_{\infty}}$$

When hiding is complete, the reflectance of the colorant doesn't depend on the concentration.

• Until when critical concentration is reached, where the assumptions of K-M model break.





Surface Reflection

Considering Surface Reflection

So far we have ignored reflection and refraction at the air/medium interface.

• Note that reflection and refraction are just macroscopic behaviors of light scattering (with interference) off of surface particles. We can reason about it using geometric optics, since a colorant is large enough for that.

Some lights are reflected right away

Air


Colorant (medium+particle)

Substrate (or another layer)

Only refracted lights undergo scattering/absorption with particles under surface


Reflectance = Surface Reflection + Back Scattering

The overall reflection is made up of both surface reflection (photons turned away right at the interface) and back-scattering from photon-particle interactions under the surface (usually called **sub-surface scattering**).




K-M Model With Surface Reflection Correction

This is called the **Saunderson correction**



K-M Model With Surface Reflection Correction



Using Fresnel equation

- r_s is 0.092 for diffuse light and 0.04 for when incident lights are perpendicular to the surface.
- r_i is 0.06 for when the internal lights crossing the interface are diffuse.

In practice, parameters are fit with observations, just like other parameters in K-M model.

Material Classification

In graphics, material is mainly classified into two categories.

- Those that can modeled by considering only surface reflection: absorption efficiency of sub-surface particles is so high that back-scattering contributes little to reflectance.
- Those where sub-surface scattering must be modeled. In graphics, people say:
 - "modeling translucent material must consider sub-surface scattering"
- "sub-surface scattering is caused by photons exiting at a point different from the incident point"

A big part of photorealistic rendering is material modeling. It's always a tradeoff between rendering time vs. realism.

But note that you don't have to be physically correct, just need to be photorealistic.



Modeling Surface


What's this type?

Surface reflection is modeled by the "roughness" of the surface.

• Rough surfaces can be modeled as a collection of small mirrors or "microfacets".

Surface types:


- Purely reflective. No roughness; perfect mirror.
- Glossy (shiny). Microfacet orientations are similar but not the same.
- Diffuse (matte). Lights reflected multiple times between "micro-mirrors". Result: reflecting evenly to all angles regardless incident angle.



Surface Types


Mirror/Glass (Specular)






Glossy (Specular)






Diffuse (Not Specular)





Puzzle






















Titanium-rutile crystal & Rutile pigment

Layering vs. Mixing

Layering




$$R_{1} = \frac{1 - [R_{g}]a_{1} - b_{1} \coth(b_{1}S_{1}X_{1})]}{a_{1} - [R_{g}] + b_{1} \coth(b_{1}S_{1}X_{1})}$$

$$R_{2} = \frac{1}{a_{2} + b_{2} \coth(b_{2}S_{2}X_{2})}$$

* Ignoring Saunderson correction here

Mixing


$$R_X = \frac{1 - R_g[a - b \coth(bSX)]}{a - R_g + b \coth(bSX)} \qquad a = \frac{K + S}{S} \qquad b = \sqrt{a^2 - 1} \qquad \coth x = \frac{e^{2x} + 1}{e^{2x} - 1}$$



$$K = \sum_i K_i c_i$$
 Relative concentration of each pigment $S = \sum_i S_i c_i$ Absorption and scattering coefficients of each pigment

Mixing When Hiding is Complete

$$\frac{K}{S} = \frac{\sum_{i} K_{i} c_{i}}{\sum_{i} S_{i} c_{i}} = \frac{(1 - R_{\infty})^{2}}{2R_{\infty}}$$



This set-up is commonly used to determine the K/S coefficients of different pigment.

- Based on reflectance measurement.
- See this paper.

K/S can also be theoretically derived, but since K-M is an empirical approximation, fitting observed data is better.

Layering == Mixing For Transparent Materials

Transparent materials do not scatter light and only absorb lights. They only take photons out.


It's not important whether the pigments are mixed or layered, i.e., where the pigment particles are. What matters is just the total amount of particles.


assuming that the absorption behaviors of individual particles are independent.


Light Colors are Additive


Mixing lights:


- If we have three lights whose *linear* RGB values are $[R_1, G_1, B_1]$, $[R_2, G_2, B_2]$, and $[R_3, G_3, B_3]$, when we mix them the mixed light's color is $[R_1 + R_2 + R_3, G_1 + G_2 + G_3, B_1 + B_2 + B_3]$.
- Can you prove it?
- The key is that the spectral power distribution of the mixed light is the sum of that of each light. From SPD to RGB color is linear.














Subtractive color space applies ONLY to strictly transparent material.


- Nearly perfect for slides/transparencies printed with dyes, pretty good for traditional photographic prints on photographic media.
- Rather poor for today's digital color printers, because they also scatter lights.












Is Mixing Materials Subtractive?

Conventional wisdom is that subtractive color model should be used when discussing paint/material mixing. Really?

Cyan + Yellow = Green because:

- Cyan material absorbs Red light; Yellow material absorbs Blue light
- Cyan and Yellow mixture would then absorb both Red and Blue, leaving only Green

By this logic, Blue + Yellow = Black/Gray





Mixing Materials is (Generally) NOT Subtractive!

Mixing paints: does subtractive color space apply? A few issues.

- First of all, a paint's color depends on illumination. So it's meaningless to discuss the color of a paint (or any non-emitting material) without illumination information.
- But more importantly paints also scatter lights.

To calculate the color of a paint mixture, we should:

- Get the spectral scattering and absorption coefficient for each pigment
- Calculate the reflectance spectrum of the mixture using the K-M model
- Use the spectrum of the illumination to calculate the R, G, B values of the mixture

True or False?





Figure 3.7: [Schematics.] Subtractive color combination. (a) A "yellow" filter transmits a band of wavelengths including spectral green, yellow, and red. It blocks other wavelengths, including those corresponding to blue and cyan. (b) Similarly, a "cyan" filter instead blocks red and yellow light. (c) Placing these two filters in succession blocks everything except green light. (d,e,f) Yellow and cyan chromophores are distributed within the fibers of a piece of paper. Multiple scattering of light combines their effects, similarly to (c).

Artists Know Better



The RYB model used by artists






Add White to Adjust Saturation Levels




Skillful painters usually use just three primary paints plus white to create different saturation levels




Masstone vs. Tint





Masstone: pure color

Tint: mixing pure color with white (desaturation)



Applicability of K-M Model

A precise theoretical material model would require modeling of both the 3D microstructure and the non-isotropic nature of the flux. However, we will almost never have the data required for such modeling and therefore have to rely on simplifying assumptions and often empirical modifications, which is what K-M does.

The accuracy is determined by the validity of the simplifications and how well the empirical modifications are able to fit the data. At some point, the empirical modifications are more like parametric functions to **fit observations** rather than rigorously justified physical models.

Painting vs. Dyeing vs. Printing

Painting vs. Dyeing vs. Printing

Similarities:

- Mixing colorants to produce color
- All have a substrate (e.g., canvas vs. textile vs. paper) on which colorants are applied

The Kubelka-Munk model has been applied to all three areas for predicting colors, reverse-engineering colorants given a sample, etc.






Colorant Differences

Painting uses **pigments**, and dyeing uses **dyes**.

- Dye molecules are orders of magnitude smaller than pigment particles.
- Dyes are water soluble (chemically bond with the water molecules).
- So dyes scattering efficiency is quite low, mostly transparent/translucent.

Printing inks can either be pigments or dyes.

Colorant Differences





Layering vs. Mixing

Painting can be layering or mixing depending on what artists decide to do.

- Glazing/vanishing are translucent layers.
- Pentimento is intentionally painting over/hiding under-layers.


Dyeing colorants are usually (pre-)mixed.

Printing inks are complicated; they are neither completely layered nor completely mixed.

Substrate Interaction

Painting: paint usually completely hides the substrate so the color usually doesn't depend on the substrate.

Dyeing: dyes are molecularly dispersed into the substrate (fibers). Dyes, while absorb lights like crazy, do not scatter nearly as much as the fibers, which contribute to most of the scattering and participate in color formation.





Substrate Interaction

Printing: somewhat in-between.

- The ink-mixture penetrates into the substrate (paper), but doesn't completely absorb lights (you can still see light from the other side!)
- So the paper contributes to color formation and is effectively just another colorant in the K-M model.

Cross sections of Epson Ultrachrome 7 pigmentbased ink printed on paper. Pigment particles are larger. They are not fully penetrated. Cross sections of Epson R340 dye-based ink printed on paper. Dye particles are smaller; they penetrate into paper well.

